

Article type : Consensus Statement

Position Statement by The Pelvic Floor Society on behalf of The Association of Coloproctology of Great Britain and Ireland on the use of mesh in ventral mesh rectopexy (VMR).

MA Mercer-Jones MD^{1#}, SR Brown MD², CH Knowles PhD³, AB Williams MD⁴

¹Consultant Colorectal Surgeon, Queen Elizabeth Hospital, Gateshead NHS Trust

²Professor of Surgery, Sheffield Teaching Hospitals, Sheffield, UK.

³Professor of Surgery, National Bowel Research Centre, Blizard Institute, Queen Mary University London, London, UK.

⁴Consultant Colorectal and General Surgeon, Guy's and St Thomas' NHS Foundation Trust, London, UK.

#Corresponding author: mark.mercer-jones@ghnt.nhs.uk

Executive summary

The following position statement forms part of a response to the current concerns regarding use of mesh to perform rectal prolapse surgery. It highlights the actions being pursued by The Pelvic Floor Society (TPFS) regarding clinical governance in relation to ventral mesh rectopexy (VMR). The following are summary recommendations:

- Available evidence suggests that mesh morbidity for VMR is far lower than that seen in transvaginal procedures (the main subject of current concern) and lower than that observed following other abdominopelvic procedures for urogenital prolapse e.g. laparoscopic sacrocolpopexy;

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/codi.13893

This article is protected by copyright. All rights reserved.

- Ventral mesh rectopexy (VMR) should be performed by adequately trained surgeons who work within a multidisciplinary team framework. Within this, it is mandatory to discuss all patients considered for surgery at a MDT meeting.
- Clinical outcomes of surgery and any complications resulting from surgery should be recorded in the TPFS-hosted national database (registry) available for this purpose; in addition, all patients should be considered for entry into ongoing and planned UK / European randomized studies where this is feasible;
- A move towards accreditation of UK units performing VMR will improve performance and outcomes in the long-term;
- An enhanced program of training including staged porcine, cadaveric and preceptorship will ensure competency of surgeons undertaking VMR;
- Enhanced consent forms and patient information booklets are being developed and these will help both surgeons and patients;
- There is weak observational evidence that technical aspects of the procedure can be optimized to reduce morbidity rates. Suture material choice may contribute towards morbidity. The available evidence is insufficient to support the use of one mesh over another (biologic versus synthetic), however the use of polyester mesh is associated with increased morbidity.

The Pelvic Floor Society

The Pelvic Floor Society (TPFS) is an affiliate of ACPGBI, and was formed to set and raise standards in pelvic floor surgery and to update members on the advances of the specialty. TPFS has a diverse membership including colorectal surgeons, urogynaecologists, urologists, radiologists, gastroenterologists, chronic pain practitioners, specialist nurses, physiotherapists and GI physiologists. The society collaborates with other pelvic floor organisations including The United Kingdom Continence Society and The British Society of Urogynaecology.

As a responsible society, and in light of ongoing concerns by the media and public groups surrounding the use of mesh in patients with pelvic organ prolapse (POP) and female stress urinary incontinence (SUI), this statement aims to address the use of mesh (synthetic and biologic) in VMR using current available evidence. Further impetus for writing this statement is in response to an ACPGBI Delphi process which ranked VMR outcomes and mesh choice as number 8 in the list of important non-cancer related questions (1). In particular the subjects of patient selection, mesh

morbidity, efficacy, clinical governance and research and development will be examined. This is not intended to be an exhaustive systematic review, nor is it a formal Delphi or RAND style consensus. It is expert opinion heavily informed by the available evidence.

Background

VMR using autologous graft, and then synthetic mesh, was first reported in 1972 by Orr-Loygue (2) for the treatment of external rectal prolapse (ERP). Unlike posterior rectopexy where simple sutures may be employed to suspend the dissected rectum, VMR requires a mesh that is fixed to the rectum and then suspended from the sacral promontory. A laparoscopic ventral approach (termed laparoscopic VMR or LVMR) was adopted and supported by D'Hoore (3) for ERP citing reduced risk for post-operative constipation compared with the posterior approach and the ability to address co-existent middle compartment prolapse in a minimally invasive fashion. Improved knowledge and insight into the pathophysiology of obstructed defaecation syndrome (ODS) and the role of internal rectal prolapse (IRP) (usually combined with rectocele), has made LVMR a procedure of choice for this condition. Surgical take-up was rapid especially within Europe for LVMR for ERP and ODS despite a lack of high-level evidence for safety or efficacy (16).

In the USA in 2008 the FDA issued a public health notification on serious complications caused by the trans-vaginal placement of mesh in woman treated for POP and SUI (4). In 2016 the FDA issued a final order to mesh manufacturers and the public to reclassify these devices from class II (moderate-risk) to class III (high-risk). The use of trans-vaginal mesh in the USA has fallen by 40-60% since (5). In Scotland in 2014, a group of women affected by mesh-related complications (mesh survivors campaign) gave evidence to the Holyrood Petitions Committee. As a result the Scottish minister for health wrote to Scottish health boards requesting a suspension of the use of mesh for POP and SUI pending an official enquiry. In 2017 the Scottish Independent review of the use, safety, and efficacy of trans-vaginal mesh implants in the treatment of POP and SUI in women was published (6). In the conclusions the Chairman stated that:

- 1) there should be patient-centred care with adequate patient choice and shared decision making supported by robust clinical governance arrangements involving a multi-disciplinary team (MDT) approach;

- 2) evidence of working in a MDT together with audit and the mandatory recording and reporting of adverse events should be a necessary part of a consultants appraisal and revalidation;
- 3) there should be informed consent;
- 4) recording databases should be improved to a national level with the creation of new data codes that would allow better NHS capture;
- 5) trans-vaginal meshes should not be offered routinely.

In the UK in 2014, the Medicines and Healthcare Products Regulatory Agency (MHRA) also published its conclusions on the evidence of the benefits and safety of vaginal mesh implants (7). These concluded that for the majority of women vaginal mesh implants were safe and effective, and when used correctly they could help alleviate POP and SUI. In addition the benefits of mesh outweighed the risks. Contrary to these conclusions, the European Commission published a report lead by its Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR) in 2015 (8). It stated that the implantation of any mesh via the vaginal route should only be considered in complex cases, in particular after a failed primary repair. Furthermore, SCENIHR opined that mesh use in SUI was acceptable if performed for moderate/severe SUI by experienced surgeons. A recent Cochrane Review in 2016 (9) on trans-vaginal mesh concurred with the SCENIHR conclusions.

The use of mesh in VMR is not analogous to trans-vaginally placed mesh. It does however have similarities with mesh use in trans-abdominal sacrocolpopexy performed for vaginal vault prolapse. Indeed VMR is quite correctly combined with sacrocolpopexy by some colorectal surgeons for single stage correction of multi-compartment prolapse. The safety and efficacy of mesh use for sacrocolpopexy was not called into question in recent enquiries, however long-term mesh/suture erosion rates of 10.5% have been estimated (10). Current NICE evidence on the safety and efficacy of sacrocolpopexy using mesh appears adequate provided that normal arrangements are in place for clinical governance and audit (IPG283). There are no NICE guidelines for VMR.

In 2014, TPFS Chairman wrote to Society members suggesting that all VMR interventions should be the subject of ongoing, continuous audit and that decision-making should be part of a MDM process. In addition device-related adverse events should be reported to the MHRA.

Patient Selection for VMR

Since VMR is performed almost exclusively by a laparoscopic approach (LVMR), the following text is principally focused on this variant of the procedure.

In 2014 a consensus report on LVMR was published from a panel of international experts (11). Definitive recommendations were for ERP in patients deemed fit for general anaesthesia. Relative indications were symptomatic high-grade IRP and patients with solitary rectal ulcer syndrome, all of whom had failed maximal conservative management. Relative contraindications included male patients with IRP, morbid obesity, previous pelvic radiotherapy, high-grade endometriosis and previous sigmoid peri-diverticulitis. Absolute contraindications were pregnancy, no demonstrable pelvic anatomical problem, a hostile abdomen, severe proctitis, psychological instability and anismus resistant to conventional treatment. It was suggested that patients with multi-compartment prolapse and those with ODS/IRP be discussed at an MDT meeting.

In 2016 the results of a nationwide pelvic floor census were published based upon research undertaken by TPFS in 2014 (12). One of the notable conclusions was that complex pelvic floor surgery (including VMR) was being undertaken in some units that did not profess an interest in pelvic floor surgery and that some units did not have access to an MDT meeting. The main value of the MDT meeting is not only in the avoidance of missing any factors relevant to the decision-making process and defining the most appropriate treatment for an individual patient but also in preventing clinicians with an over enthusiastic belief in their own favoured treatment modality e.g., VMR, pursuing an inappropriate course of treatment. It also allows a robust audit process to be developed with adequate data collection.

The monitoring of standards and quality of MDT meetings and of data collection will in the future form part of UK pelvic floor unit's accreditation. This initially voluntary process is being driven by TPFS to define and monitor standards of care, organisation and quality and should reassure patients, commissioners and providers that selection, efficacy and safety are paramount in delivering care for patients undergoing VMR.

LVMR: technical considerations

Since VMR is performed almost exclusively by a laparoscopic approach (LVMR), the following text is principally focused on this variant of the procedure.

Mesh

Meshes currently used for LVMR are either synthetic or biologic. Synthetic meshes are non-absorbable, partially absorbable or absorbable. Biologic meshes are porcine-derived and either cross-linked or non-cross-linked.

The use of absorbable and semi-absorbable synthetic meshes is not advised because of concerns over recurrence rates (11). The commonest non-absorbable synthetic meshes in use are, polypropylene and polyester. Polypropylene can be light or heavy-weight, and coated or uncoated.

The data regarding mesh complications and recurrence rates are summarized below for the interested reader. Critical to the interpretation of these data is a prior understanding that all current data are observational and nearly all derive from low quality studies (level IV case series or poor-quality cohort studies) or synthesis thereof. Thus, while the data have been provided, it is TPFS view that no current evidence-based recommendation can be made regarding mesh selection (in particular, the biologic vs. synthetic mesh argument).

The largest study examining outcome after LVMR (for a mix of internal and external rectal prolapse) was an observation study conducted in the Netherlands and Belgium (13). Synthetic mesh (polyethylene or polypropylene) was used in 919 patients with a mix of internal and external rectal prolapse. Median follow-up was 33.9 months (0.4-143.6). Estimated mesh-related complications assessed by survival curves and analysis (Kaplan- Meier) at 3, 5 and 10 years were 1.5%, 2.9%, and 4.6%. However, this estimation included detachment of mesh from the sacral promontory (2.7%). A multicentre collaboration to assess the safety of laparoscopic ventral rectopexy examined mesh type and complication rates for various synthetic (n=1764) and biologic (n=439) meshes implanted in over 2200 patients (14). The synthetic meshes compared were: polypropylene (0-85 g/m²), polyester, and titanium-coated polypropylene (35 g/m²); biologics were either cross-linked porcine dermis or

uncross-linked porcine intestinal submucosa. The synthetic erosion rate was 2.4% (mean FU 38 months [0-162]), and the biologic erosion rate was 0.7% (mean FU 26 months [0-68]). Kaplan-Meier estimates of erosion probability at 1, 2 and 5 years for synthetic mesh were 0.4%, 1.1%, and 2.3%. For biologic they were 0.5%, 0.7%, and 0.7%. There was no statistical difference between synthetics or biologics. However, no patients with a biologic complication required major surgical intervention compared with 40% of synthetic mesh complications. Polyester mesh was associated with a statistically-significant increase risk of erosion compared to the other mesh types.

These low rates of complications are similar to a recent systematic review comparing synthetic and biologic meshes (14). In this study, the synthetic mesh erosion rate was 1.87% (66/3517) and the biologic was 0.22% (1/439).

As part of the National Institute of Health Research (£2M) funded Constipation treatment pathway program of studies (CapaCITY), a systematic review of surgical treatments for constipation has been recently conducted for 5 main procedure approaches. One of these 5 methodologically-robust reviews addressed outcomes of forms of rectal suspension procedures (rectopexy) with LVMR being one of the main procedures considered (predominant available data). Following data synthesis of 18 eligible studies, mesh-related complications were 0.5% (0-3.9%)(16) at a median follow up of 25 months.

The above data compare highly favourably with erosion rates following sacrocolpopexy. In a systematic review these were 0-21% (17), and in the CARE trial (9), Kaplan-Meier estimate at 7 years was 10.5% (95% CI 6.8% to 16.1%). Published evidence suggests that LVMR has far less mesh-related morbidity compared with sacrocolpopexy.

Sutures

It has been postulated that some mesh-related morbidity is related to suture type. A historical cohort study compared suture type in sacrocolpopexy and mesh/suture complications (18). Patients all had polypropylene mesh implanted. Erosion rates were 3.7% for polyester (ethibond) sutures (6/161) and 0% for polydioxanone sulphate (PDS) sutures (0/254). A consensus statement expressed

a view that vaginal sutures for LVMR should be PDS (for any mesh type) and that rectal sutures should be PDS if synthetic mesh is used (11). In an effort to avoid sutures altogether, two centres have promoted the use of synthetic mesh with tissue glue (cyanoacrylate or synthetic hydrogel) to secure mesh to the rectum (19, 20). Results appear encouraging and these warrant further research.

Consent

Regarding consent for LVMR, it is imperative the operating surgeon explains that a mesh will be used and warns the patient of potential complications. Ideally risk should be based upon the surgeon's own data (a further argument for unit data submission) however registry and systematic review data are also acceptable. On the basis of the latter (and uncertainty due to poor data quality and study design) it is reasonable to quote an overall mesh complication rate of 2.5% based on worst-case scenario. These figures will potentially be higher for re-do surgery. Patients should also be warned of the rare risks of discitis from fixation to the sacral promontory. TPFS is producing an enhanced LVMR consent form together with patient information documentation. This will also integrate patient's views on language based on a national meeting (Birmingham June 2017).

Efficacy of LVMR

Evidence for efficacy for LVMR either for ERP or ODS/IRP is of poor quality. Excepting one study, there are no randomized trials examining outcome for LVMR. This small RCT was a double-blind, randomized, single centre study comparing laparoscopic posterior suture rectopexy (n=37) with LVMR using polypropylene mesh (n=38) for patients with ERP (21). The primary outcome was a change in the ODS score at 12 months. Both procedures resulted in significant reductions in the ODS score (mean 2.18 [95% CI -0.14 to 4.49]) with no significant difference in effect size between groups. There was no significant difference in recurrence rates (suture 5%, LVMR 0%), and no mesh related morbidity. A further trial is seeking NIHR funding to address which is the superior procedure with validated PROMs and scoring systems, and longer follow-up (for recurrence and mesh morbidity).

Observational data include a variety of cohort designs of varying quality (mostly level IV evidence). The study with the longest follow-up by D'Hoore (13), included 919 patients with ODS (n=677)_(with IRP and/or symptomatic rectocele) and patients with ERP (n=242). Improvement in faecal incontinence was seen in 80% of all patients and improvement in ODS scores in 74% of those with

ODS, and 61% of those with ERP. Estimated overall prolapse recurrence rates estimated by Kaplan-Meier at 3, 5, and 10 years were 7%, 10.7%, and 14.3%. The recently conducted systematic review conducted as part of the NIHR CapaCITY program provides the following synthesized data estimates and graded practice recommendations:

- For rectal suspension procedures performed for ODS (the majority being LVMR), global satisfaction was 83% (74-91%) (level of evidence IV, graded practice recommendation C)(16). For LVMR, there was an 86% improvement in constipation (level of evidence IV, graded practice recommendation C). Data were predominately derived from level IV observational studies and it is known that observational bias can affect results.
- There were no data comparing efficacy of biologic versus synthetic mesh use in LVMR, and any outcome data for biologic mesh is short-term only.

Proficiency

There is no doubt that LVMR can be challenging technically. If done incorrectly there is not only a risk to efficacy but also potential for harm. There is undoubtedly a learning curve to achieve proficiency. Although this is difficult to estimate, the adoption of CUSUM curve methodology suggests in terms of operating time this learning curve takes approximately 54 cases before stability is achieved (22). Such numbers are difficult to achieve even in a high volume centre. However, it is possible to reduce the learning curve with robust mentorship and training (23). If a rigorous training programme is undertaken (including theory assessment and simulation), estimates of the learning curve using the same methodology suggest proficiency can be achieved after around 25-30 cases.

In order to provide this rigorous training, TPFS has been engaged in devising a specific curriculum, developing hands on training courses, providing mentorship and organising regular education events where issues can be discussed in an open forum with experienced surgeons. The Pelvic Floor Society website (24) contains various teaching aids including operative videos. Specific pelvic floor fellowships are encouraged and advertised through this website.

Accreditation

It is essential to identify UK units that have the necessary expertise and infrastructure to deliver high quality including surgical expertise. A recent census identified 67 centres who provide a pelvic floor service (25). In 39% of these centres there was significant infrastructure and support for pelvic floor

Accepted Article

services and the volume of LVMR procedures was high (median 20 per annum). These units were often tertiary referral centres for their region. Other centres had less support but still delivered a median of 12 cases per year. In 13% there was little specific pelvic floor interest but nevertheless a median of 3 procedures occurred per year. Whilst volume does not necessarily equate to optimal outcome (26) it is essential that there is a process to assess all centres whatever their volume of work. Accreditation is one method of assessment. If carried out correctly accreditation not only allows identification of units providing high quality care but introduces a mindset of high performance bringing up below standard units and improving further those who meet the set standards (27). This benefits the patient first and foremost in terms of confidence, choice and influence. However there are also benefits to health professionals in terms of recognition and leverage, to Trusts in terms of quality assurance and to regulators and commissioners in terms of risk reduction and value for money. A part of the accreditation process will be the assessment of a pelvic floor unit's MDT approach. TPFS robustly endorse the MDT approach. As in the Scottish independent review (6) on recommendations for urogynaecology units performing vaginal prolapse surgery, for pelvic floor units performing VMR, there should be patient-centred care with adequate patient choice and shared decision making supported by robust clinical governance arrangements involving a MDT approach. In addition evidence of working in a MDT together with audit and the mandatory recording and reporting of adverse events should be a necessary part of a consultants appraisal and revalidation.

TPFS is in the process of developing a voluntary accreditation process along the lines of that already in practice in Urogynaecology (28).

The need for expertise is nowhere more acute than in the care of patients who have had mesh-related complications. Such patients may require very complex surgery with a significant incidence of morbidity. This is also true for surgery for recurrent prolapse. There is evidence that suggests timely and appropriate intervention in these groups can reduce the potential morbidity and allow restoration of normal function (29). TPFS has identified UK units that profess to have this expertise (24) and surgeons are encouraged to refer to these teams early if complications occur.

Research and development

Many surgeons are convinced by the perceived benefits of a LVMR in terms of reduced symptom recurrence and improved functional outcome compared with alternative therapies. The procedure is now the most commonly performed abdominal procedure in the UK for full thickness rectal prolapse

(30). Nevertheless, the evidence base remains weak (31, 32, 33). There is a desperate need for high quality UK audit data and research in this area. TPFS has developed and introduced a carefully designed database (33) that will allow assessment of current practice. Data entry is, however, voluntary, but all surgeons that carry out this procedure are encouraged to enter their cases into this database. In addition, any adverse incident involving a mesh should be reported to the MHRA (35).

Regarding research, two relevant UK initiatives merit mention. The first, CapaCiTY 3 (16) is an NIHR-funded randomised trial that will evaluate efficacy for LVMR in patients with ODS using a stepped-wedge design. This study is currently recruiting in 8 UK centres. Further research into full thickness rectal prolapse is planned with a grant application submitted based on recent quality evidence from Denmark comparing posterior suture rectopexy with LVMR (21).

Given the current concerns regarding mesh related harm all surgeons that utilise this procedure should be encouraged to offer each and every patient the opportunity to be part of such a trial.

Patient Commentary

Whilst all surgery is not without risk of complications, a small number of patients have reported distressing symptoms following pelvic surgery where mesh implants have been used. In addition, various international regulatory bodies have disagreed with regard to the risk status of using such implants, as well as a lack of clarity with regard to the extent of risk for each of the different types of operation where mesh is used to treat prolapse of the various different organs which make up the pelvic floor.

Patients should know that where surgery is used to correct external rectal prolapse or troublesome internal prolapse, there is a lower risk of mesh failure than with surgery to correct the weakness of the other pelvic organs where mesh is used. This is because the mesh for the VMR operation is inserted between the rectum and the vagina rather than directly into the vagina.

Patients should be aware that VMR, with which this Statement is concerned, is the best available treatment in the UK to restore normal rectal function. This remains the case until any fresh evidence may arise.

Accepted Article

Patients present and future can be reassured that all measures are underway to ensure that all aspects of delivering a safer service are being explored and recorded in order to understand why a small number of patients are affected, and what can be done to make this procedure as safe as possible in the future. In this way, it is hoped that an eventual “gold standard” service in VMR will be achieved for patients.

New guidelines have been provided to surgeons with greater attention to further training and a database is already underway, which will give information on patient outcomes. The various surgical specialists who deal with pelvic floor problems are working together to pool their expertise to try to understand this complex area of surgery, and to refine surgical techniques. All patients should now benefit from the combined opinions of a range of experts in the pelvic floor when their clinical case is discussed before treatment. Hospital units, which deliver this operation, are to be examined and accredited to ensure the highest standards of care. However, the implementation and on going research will take some time, but every effort is being made to improve greater understanding of this area of surgery and therefore how to minimise complications.

Importantly, measures are now being taken to prepare improved patient information to help patients in the discussion they have with their surgeon with regard to weighing up the risks of treatment against the benefits of a potential return to improved rectal function, where other treatments have failed. Patients may wish to explore with their surgeon their individual risk profile, with an understanding of the type and extent of the prolapse for which treatment is sought, and to ensure that they are also fully prepared for the slim chance of complications.

REFERENCES

1. Tiernan J, Cook A, Geh I, George B, Magill L, Northover J, Verjee A, Wheeler J, Fearnhead N. Use of a modified Delphi approach to develop research priorities for the association of coloproctology of Great Britain and Ireland. *Colorectal Dis.* 2014 Dec;16(12):965-70.
2. Loygue J, Huguier M, Malafosse M, Biotois H. Complete prolapse of the rectum. A report on 140 cases treated by rectopexy. *Br J Surg.*1971 Nov;58(11):847-8.
3. D’Hoore A, Cadoni R, Penninckx F. Long-term outcome of laparoscopic ventral rectopexy for total rectal prolapse. *Br J Surg.* 2004 Nov;91(11):1500-5.
4. <https://www.fda.gov/downloads/medicaldevices/safety/alertsandnotices/ucm26276.pdf>

- Accepted Article
5. Daly J, Frazer M. Decline and Fall, lessons learned from the troubled history of transvaginal mesh kits. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists; O&G Magazine 2014; 16(1), 2014.
 6. The Scottish Independent Review of the Use, Safety and Efficacy of Transvaginal Mesh Implants in the Treatment of Stress Urinary Incontinence and Pelvic Organ Prolapse in Women: Final Report www.gov.scot/Publications/2017/03/3336.
 7. ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_049.pdf
 8. https://ec.europa.eu/health/sites/health/files/scientific_committees/emerging/docs/scenih_r_o_049.pdf
 9. Maher C, Feiner B, Baessler K, Christmann-Schmid C, Haya N, Marjoribanks J. Transvaginal mesh or grafts compared with native tissue repair for vaginal prolapse. Cochrane Database of Systematic Reviews 2016;2: CD012079
 10. Nygaard I, Brubaker L, Zyczynski HM et al. Long-term outcomes following abdominal sacrocolpopexy for pelvic organ prolapse. JAMA 2013 May;309(19):2016-24.
 11. Mercer-Jones MA, D'Hoore A, Dixon AR, Lehur P, Lindsey I, Mellgren A, Stevenson AR. Consensus on ventral rectopexy: report of a panel of experts. Colorectal Dis. 2014 Feb;16(2):82-8.
 12. Hainsworth AJ, Schizas AM, Brown S, Williams AB. The future of pelvic floor services in the UK. Colorectal Dis. 2016 Nov;18(11):1087-1093.
 13. Consten EC, van Iersel JJ, Verheijen PM, Broeders IA, Wolthuis AM, D'Hoore A. Long-term Outcome After Laparoscopic Ventral Mesh Rectopexy: An Observational Study of 919 Consecutive Patients. Ann Surg. 2015 Nov;262(5):742-7
 14. Evans C, Stevenson AR, Sileri P, Mercer-Jones MA, Dixon AR, Cunningham C, Jones OM, Lindsey I. A Multicenter Collaboration to Assess the Safety of Laparoscopic Ventral Rectopexy. Dis Colon Rectum. 2015 Aug;58(8):799-807
 15. Balla A, Quaresima S, Smolarek S, Shalaby M, Missori G, Sileri P. Synthetic versus biologic mesh-related erosion after ventral mesh rectopexy. A systematic review. Ann Coloproctol 2017 April;33(2):46-51.
 16. Brown SR, Grossi U, Lacy-Colson J, Mercer-Jones M, Mason J, Knowles CH. SURGERY FOR CONSTIPATION: SYSTEMATIC REVIEW AND PRACTICE RECOMMENDATIONS. RESULTS II: HITCHING PROCEDURES FOR THE RECTUM (RECTAL SUSPENSION) In Press Colorect Dis 2017.

- Accepted Article
17. Jia X, Glazener C, Mowatt G, Jenkinson D, Fraser C, Bain C, Burr J. Systematic review of the efficacy and safety of mesh in surgery for uterine or vaginal vault prolapse. *Int Urogynecol J*. 2010 Nov;21(21-22):1413-31.
 18. Shepherd JP, Higdon HL 3rd, Stanford EJ, Mattox TF. Effect of suture selection on the rate of suture or mesh erosion and surgery failure in abdominal sacrocolpopexy. *Female Pelvic Med Reconstr Surg*. 2010 Jul;16(4):229-33.
 19. Silveira RK, Domingie S, Kirzin S, de Melo Filho DA, Portier G. Comparative study of safety and efficacy of synthetic surgical glue for mesh fixation in ventral rectopexy. *Surg Endosc*. 2017 Mar 31. [Epub ahead of print]
 20. Gravie JF, Magnie C. "Wrap technique": a new operative procedure using a self-adhesive prosthesis for laparoscopic ventral rectopexy. *Tech Coloproctol* 2015 June;19(6):361-3.
 21. Lundby L, Iverson LH, Bunzen S, Wara P, Hoyer K, Laurberg S. Bowel function after laparoscopic posterior sutured rectopexy versus ventral mesh rectopexy for rectal prolapse: a double-blind, randomised single-centre study. *Lancet Gastroenterol Hepatol*. 2016 Dec;1(4):291-297
 22. Mackenzie H, Dixon AR. Proficiency gain curve and predictors of outcome for laparoscopic ventral mesh rectopexy. *Surgery*. 2014 Jul;156(1):158-67.
 23. Pucher PH, Mayo D, Dixon AR, Clarke A, Lamparelli MJ, Learning curves and surgical outcomes for proctored adoption of laparoscopic ventral mesh rectopexy: cumulative sum curve analysis. *Surg Endosc*. 2017 Mar;31(3):1421-1426.
 24. thepelvicfloorsociety.co.uk
 25. <https://www.acpgbi.org.uk/news/resources-for-coloproctology-20150>
 26. Faiz O. The volume-outcome relationship in colorectal surgery. *Tech Coloproctol*. 2014 Oct;18(10):961-2.
 27. Greenfield D, Kellner A, Townsend K, Wilkinson A, Lawrence SA. Health service accreditation reinforces a mindset of high-performance human resource management: lessons from an Australian study. *Int J Qual Health Care*. 2014 Aug;26(4):372-7
 28. <http://bsug.org.uk/pages/information/accreditation-of-units/102>
 29. Badrek-Al Amoudi AH, Greenslade GL, Dixon AR. How to deal with complications after laparoscopic ventral mesh rectopexy: lessons learnt from a tertiary referral centre. *Colorectal Dis*. 2013 Jun;15(6):707-12.
 30. Gunner CK, Senapati A, Northover JM, Brown SR. Life after PROSPER. What do people do for external rectal prolapse? *Colorectal Dis*. 2016 Aug;18(8):811-4.

- Accepted Article
31. Gouvas N, Georgiou PA, Agalianos C, Tan E, Tekkis P, Dervenis C, Xynos E. Ventral colpoproctopexy for overt rectal prolapse and obstructed defaecation syndrome: a systematic review. *Colorectal Dis.* 2015;17(2):O34-46.
 32. Murphy PB, Wanis K, Schlachta CM, Alkhamesi NA. Systematic review on recent advances in the surgical management of rectal prolapse. *Minerva Chir* 2017 Feb;72(1):71-80.
 33. Tou S, Brown SR, Nelson R. Surgery for complete (full-thickness) rectal prolapse in adults. *Cochrane Database Syst Rev.* 2015;11:CD001758.
 34. https://npfs.herokuapp.com/users/sign_in
 35. <https://www.gov.uk/government/organisations/medicines-and-healthcare-products-regulatory-agency>

DISCLOSURES

- CHK is a paid consultant and speaker for Medtronic in relation to sacral neuromodulation (no involvement with mesh or rectopexy instrumentation)
- MMJ is a preceptor for Medtronic in relation to LVMR.
- ABW is a non paid consultant for Cook Medical and Medtronic in relation to pelvic floor surgery and anal fistula surgery

Acknowledgements.

We are grateful to the following members of TPFS executive committee for reviews of earlier drafts of this manuscript, all have seen the final version and are in agreement for its submission: Jane Dixon, Jennie Grainger, Oliver Jones, Simon Phillips, Bandipalyam Praveen, Jon Randall, Shahab Siddiqi, Karen Telford, Dorin Ziyaie. In addition we are grateful to Jo Church, Chair of ACPGBI Patient Liaison Group for producing the patient commentary.